

ARE ABSOLUTE ABUNDANCES RELATED TO TARSAL PAD LENGTH IN *CENTROBOLUS* COOK, 1897?

Mark Cooper*

University of South Africa, South Africa.

*Correspondence email, cm.i@aol.com, +27714620070.

Abstract- Two species of *Centrobolus* were identified (*C. anulatus*, *C. inscriptus*) based on morphology and confirmed using Scanning Electron Microscopy (SEM) of gonopod and tarsal pads. Three sets of measurements were made from data: (1) tarsal pad length (μm), (2) tarsal pad to tarsus length ratios, and (3) absolute abundances. Absolute abundances and tarsal pad lengths were correlated ($r=0.63$, Z score= 1.66 , $n=8$, $p<0.05$). Tarsal pad lengths to tarsus ratios and absolute abundances were positively related ($r=0.63046242$, Z score= 1.65957221 , $n=8$, $p=0.04850025$). These new relationships between tarsal pad length, tarsal pad to tarsus ratios, and absolute abundances among the *Centrobolus* millipedes support the function of the tarsal pads as adaptive characters in mate competition and securing mates among increased absolute abundances.

I. INTRODUCTION

The red millipede genus *Centrobolus* is well known for studies on sexual size dimorphism (SSD) and displays prolonged copulation durations [5-10]. *Centrobolus* is distributed in temperate southern Africa with northern limits on the east coast of southern Africa at -17° latitude South (S) and southern limits at -35° latitude S. It consists of taxonomically important species with 12 species considered threatened and includes nine vulnerable and three endangered species [25]. It occurs in all the forests of the coastal belt from the Cape Peninsula to Beira in Mocambique [24]. Spirobolida has two pairs of legs modified into gonopods on the eighth and ninth diplosegments [27]. In *Centrobolus* the coleopods are the anterior gonopods of leg-pair eight. They can be classed as paragonopods or peltogonopods because they are fused into a single plate-like structure and play a subsidiary role as inseminating devices. In contrast, leg-pair nine is sperm-transferring [2]. The sternites (or stigma-

carrying plates [26]) prevent lateral shifting (stabilizer) and stretch the vulva sac in a medial plane [5].

These worm-like millipedes have female-biased SSD [5-10, 13-20, 22]. From the results, correlations between tarsal pad lengths and absolute abundances were checked for correlations.

II. MATERIALS AND METHODS

Millipedes were hand collected in coastal forest habitat at Mtunzini ($28^\circ 55' \text{ S}$; $31^\circ 45' \text{ E}$) during the summer season (1995-1996). Individual millipedes were identified as species and sexed based on the presence of gonopods in males and their absence in females. Individuals were counted as either on or above ground ($>30\text{cm}$ but $<3\text{m}$ above ground surface). The number of mating pairs was recorded. The total number of adults was used to estimate the relative abundance. Intercalary males were excluded from the counts. Two species of *Centrobolus* were identified based on morphology and confirmed using Scanning Electron Microscopy (SEM) of gonopod structure (*C. anulatus*, *C. inscriptus*). The tarsi and gonopods were dissected from males of these two species and prepared for SEM. Specimens were fixed, first in 2.5% glutaraldehyde (pH 7.4 phosphate-buffered saline) at 4°C for 24 hours, then in osmium tetroxide (2%). Dehydration through a graded alcohol series (50%, 60%, 70%, 80%, 90% to 100% ethanol) and critical point drying followed. Specimens were mounted on stubs and sputter coated with gold palladium. Tarsal pads and gonopods were viewed under a Cambridge S200 SEM. SEM micrographs were examined and the individual

components of the gonopods were identified according to the available species descriptions. Two sets of measurements were made from the micrographs (1) tarsal pad length, (2) ratio of tarsal pad length to total tarsus length and the third set included field data for (3) absolute abundances. Tarsal pad lengths, tarsal pad to tarsus ratios, and absolute abundances were correlated using Pearson's Correlation Coefficient (<https://www.gigacalculator.com/calculators/correlation-coefficient-calculator.php>).

III. RESULTS

Tarsal pad lengths in *C. anulatus* (540µm, n=8) and *C. inscriptus* (1018µm, n=10), tarsal pad length to tarsus ratios in *C. anulatus* (0.83), and *C. inscriptus* (1.04), and absolute abundances were given [1, 3]. Tarsal pad lengths and abundances were positively related (Figure 1: $r=0.63046242$, Z score=1.65957221, $n=8$, $p=0.04850025$). Tarsal pad lengths to tarsus ratios and abundances were positively related (Figure 2: $r=0.63046242$, Z score=1.65957221, $n=8$, $p=0.04850025$).




Figure 1. Relationship between tarsal pad length (Y) and absolute abundances (x) for *C. anulatus* and *C. inscriptus*.




Figure 2. Relationship between tarsus pad length to tarsus ratio (Y) and absolute abundances (x) for *C. anulatus* and *C. inscriptus*.

IV. DISCUSSION

The tarsal pads are important taxonomic and secondary sexual characters of *Centrobolus* [24]. The tarsal pad length, tarsal pad-to-tarsus ratios, and absolute abundances were estimated in two *Centrobolus* species [3]. A direct relationship between three factors (tarsal pad length, tarsal pad to tarsus ratios, and absolute abundances) in the millipedes is compared which certainly supports the relationship. A relationship between these behavioral and morphological factors is present across the two species suggesting an adaptive character. *C. inscriptus* had higher absolute abundances and longer tarsal pads while *C. anulatus* had lower absolute abundances and a shorter tarsal pad length and tarsal pad to tarsus ratio. There were two positive relationships between tarsal pad length, tarsal pad length to tarsus ratio, and absolute abundances.

V. CONCLUSION

New relationships between tarsal pad length, tarsal pad to tarsus ratios, and absolute abundances among the *Centrobolus* millipedes support the function of the tarsal pads as adaptive characters in mate competition and acquiring mates among increased absolute abundances.

APPENDIX.

Male and female absolute abundances (early, and late in a season, on the ground, and in the trees), in two species of *Centrobolus* followed by tarsal pad lengths (μm) and tarsal pad length to tarsus ratio.

0, 540, 0.83 (*C. anulatus*).

58, 540, 0.83 (*C. anulatus*).

101, 1018, 1.04 (*C. inscriptus*).

445, 1018, 1.04 (*C. inscriptus*).

75, 540, 0.83 (*C. anulatus*).

800, 1018, 1.04 (*C. inscriptus*).

46, 540, 0.83 (*C. anulatus*).

135, 1018, 1.04 (*C. inscriptus*).

REFERENCES

1. M. Cooper, "Sex ratios, mating frequencies and relative abundance of sympatric millipedes in the genus *Chersastus*," *Arthropods*, vol. 3, no. 4, pp. 174-176, 2014.
2. M. I. Cooper, "Confirmation of four species of *Centrobolus* Cook (Spirobolida: Trigonulidae) based on gonopod ultrastructure," *Journal of Entomology and Zoology Studies*, vol. 4, no. 4, pp. 389-391, 2016.
3. M. I. Cooper, "Tarsal pads of *Centrobolus* Cook (Spirobolida: Trigonulidae)," *Journal of Entomology and Zoology Studies*, vol. 4, no. 3, pp. 385-386, 2016.
4. M. Cooper, "Julid millipede and spirobolid millipede gonopod functional equivalents," *Journal of Entomology and Zoology Studies*, vol. 7, no. 4, pp. 333-335, 2019.
5. M. I. Cooper, "Sexual size dimorphism and corroboration of Rensch's rule in *Chersastus* millipedes," *Journal of Entomology and Zoology Studies*, vol. 2, no. 6, pp. 264-266, 2014.
6. M. I. Cooper, "Copulation and sexual size dimorphism in worm like millipedes," *Journal of Entomology and Zoology Studies*, vol. 5, no. 3, pp. 1264-1266, 2017.
7. M. Cooper, "*Centrobolus anulatus* (Attems, 1934) reversed sexual size dimorphism," *Journal of Entomology and Zoology Studies*, vol. 6, no. 4, pp. 1569-1572, 2018.
8. M. I. Cooper, "The relative sexual size dimorphism of *Centrobolus inscriptus* compared to 18 congeners," *Journal of Entomology and Zoology Studies*, vol. 4, no. 6, pp. 504-505, 2016.
9. M. I. Cooper, "Relative sexual size dimorphism in *Centrobolus fulgidus* (Lawrence) compared to 18 congeners," *Journal of Entomology and Zoology Studies*, vol. 5, no. 3, pp. 77-79, 2017.
10. M. I. Cooper, "Relative sexual size dimorphism *Centrobolus ruber* (Attems) compared to 18 congeners," *Journal of Entomology and Zoology Studies*, vol. 5 no. 3, pp. 180-182, 2017.
11. M. I. Cooper, "Competition affected by re-mating interval in a myriapod," *Journal of Entomology and Zoology Studies*, vol. 3, no. 4, pp. 77-78, 2015.
12. M. Cooper, "Re-assessment of Rensch's rule in *Centrobolus*," *Journal of Entomology and Zoology Studies*, vol. 5, no. 6, pp. 2408-1410, 2017.
13. M. I. Cooper, "Size matters in myriapod copulation," *Journal of Entomology and Zoology Studies*, vol. 5, no. 2, pp. 207-208, 2017.
14. M. I. Cooper, "Sexual size dimorphism and the rejection of Rensch's rule in Diplopoda," *Journal of Entomology and Zoology Studies*, vol. 6, no. 1, pp. 1582-1587, 2018.
15. M. I. Cooper, "Allometry for sexual dimorphism in millipedes," *Journal of Entomology and Zoology Studies*, vol. 6, no. 1, pp. 91-96, 2018.
16. M. I. Cooper, "Trigoniulid size dimorphism breaks Rensch," *Journal of Entomology and Zoology Studies*, vol. 6, no. 3, pp. 1232-1234, 2018.
17. M. Cooper, "A review of studies on the fire millipede genus *Centrobolus* (Diplopoda: Trigonulidae)," *Journal of Entomology and Zoology Studies*, vol. 6, no. 4, pp. 126-129, 2018.
18. M. Cooper, "*Centrobolus sagatinus* sexual size dimorphism based on differences in horizontal tergite widths," *Journal of Entomology and Zoology Studies*, vol. 6, no. 6, pp. 275-277, 2018.
19. M. Cooper, "*Centrobolus silvanus* dimorphism based on tergite width," *Global Journal of Zoology*, vol. 3, no. 1, pp. 003-005, 2018.
20. M. Cooper, "Xylophagous millipede surface area to volume ratios are size dependent in forest," *Arthropods*, vol. 8, no. 4, pp. 127-136, 2019.
21. J. M. Dangerfield, S. R. Telford, "Seasonal activity patterns of julid millipedes in Zimbabwe," *Journal of Tropical Ecology*, vol. 7, pp. 281-285, 1991.
22. J. M. Dangerfield, A. E. Milner, R. Matthews, "Seasonal activity patterns and behaviour of juliform millipedes in south-eastern Botswana," *Journal of Tropical Ecology*, vol. 8, no. 4, pp. 451-464, 1992.
23. M. D. Greyling, R. J. Van Aarde, S. M. Ferreira, "Seasonal changes in habitat preferences of two closely related millipede species," *African Journal of Ecology*, vol. 39, no. 1, pp. 51-58, 2001.
24. R. F. Lawrence, "The Spirobolida (Diplopoda) of the eastern half of Southern Africa*," *Annals of the Natal Museum*, vol. 18, no. 3, pp. 607-646, 1967.
25. R. P. Mailula, "Taxonomic revision and Red List assessment of the 'red millipede' genus *Centrobolus* (Spirobolida: Pachybolidae) of South Africa," *The University of Kwazulu natal*, xxiii+289, 2021.
26. P. Sierwald, J. E. Bond, "Current Status of the Myriapod Class Diplopoda (Millipedes): Taxonomic Diversity and Phylogeny," *Annual Review of Entomology*, vol. 52, no. 1, pp. 401-420, 2007.
27. X. J. Zahnle, P. Sierwald, S. Ware, J. E. Bond, "Genital morphology and the mechanics of copulation in the millipede genus *Pseudopolydesmus* (Diplopoda: Polydesmida: Polydesmidae)," *Arthropod Structure & Development*, vol. 54, pp. 1009-1013, 2020.